

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 April 2003 (10.04.2003)

PCT

(10) International Publication Number
WO 03/029440 A1

(51) International Patent Classification⁷: C12N 1/20,
C12P 7/56 // (C12N 1/20, C12R 1:225)

(21) International Application Number: PCT/SE02/01805

(22) International Filing Date: 3 October 2002 (03.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0103294-5 3 October 2001 (03.10.2001) SE

(71) Applicant (*for all designated States except US*):
FORSKARPATENT I SYD AB [SE/SE]; IDÉON,
S-223 70 LUND (SE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ANDERSSON, Ulrika** [SE/SE]; Tallgången 2, S-240 14 Veberöd (SE).
RÅDSTRÖM, Peter [SE/SE]; Lundavägen 19, S-237 37 Bjärred (SE).

(74) Agent: **STRÖM & GULLIKSSON IP AB**; Sjöporten 4,
S-417 64 Göteborg (SE).

(81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MUTATED LACTOCOCCUS STRAIN

(57) Abstract: The present invention relates to a mutant of *Lactococcus lactis* spp. *lactis*, which produces lactate at high volumetric and specific productivity as well as high amounts of lactate dehydrogenase, as well as the use of said strain, and a method for producing lactate, as well as the use of the lactate thus produced.

WO 03/029440 A1

Mutated lactococcus strain

DESCRIPTION

5 Technical field

The present invention relates to a new lactate producing strain belonging to the *Lactococcus* genus, viz. a mutant of *Lactococcus lactis* spp. *lactis* 19435 (obtained from ATCC), as well as a use of the mutant for lactate production, and a method for producing lactate.

10

Background of the invention

Lactic acid is a naturally occurring organic acid that can be produced by either chemical synthesis or carbohydrate fermentation. Both of these production routes are used commercially [Datta, (1995)]. Chemical synthesis results in racemic lactic acid, while
15 fermentation technologies enable synthesis of a desired stereoisomer of lactic acid. In the literature there are several reports concerning metabolic engineering of lactic acid bacteria in order to assess and enhance their ability in lactic acid production from different carbohydrates. Existing commercial production processes use homolactic organisms such as *Lactobacillus delbrueckii* and *Lb. bulgaricus* [Datta, (1995)].

20

Lactococcus lactis is one of the most studied organisms used for industrial applications. *L. lactis* ferments glucose according to a homolactic pathway under non-limiting glucose conditions [Sjöberg, (1995)]. This organism primarily produce the L-isomer of lactic acid that is favoured in applications associated with food since the D-isomer is harmful to
25 humans [Hofvendahl, (1997)]. The post-glycolytic pyruvate metabolic pathways play a key role in determining the outcome of a fermentation of *L. lactis*. Activities of pyruvate converting enzymes changes with different cultivation conditions which lead to the important variations of end product formation [Cocaign-Bousquet, (1996)] (and references therein). In this study we describe a mutant of *Lactococcus lactis* ssp. *lactis*
30 that is able to produce lactate twice as fast as the wild type strain under non-limiting glucose conditions. Our report concerns a physiological and biochemical characterisation of the mutant lactococcal strain in order to assess differences in glucose uptake and activities of important enzymes involved in glycolysis and pyruvate conversion compared to the wild type strain.

Lactic acid is a chemical used in food technology as well as in general chemical industry, including polymer technology. Thus, it can be used to produce polymers or become
5 hydrogenated to produce propylene glycol, and other carbon chemical intermediates. Lactic acid is of interest to produce biocompatible and decomposable polymers, polylactic acid (PLA), used to produce sutures or implants, i.e., to be used within medicinal and veterinary surgery.

10 50,000 tonnes of lactate are produced each year around the world, which indicates that it is a relatively attractive product. Two thirds are recovered from fermentation processes while one third is derived from a synthetic production from mostly lacto-nitrile. The drawback of using a synthetic production is that the lactate recovered is a racemate, i.e., containing equal amounts of L- and D-isomers in a mixture. The drawback of having a
15 racemate is that it is considerably harder to polymerise the lactate at such an application. Pure D- or L-lactate polymers are crystalline and stable, while polymers of mixtures are amorphous.

Lactic acid is an expensive chemical when produced by fermentation of different lactic
20 acid producing micro-organisms, as the fermentation produces lactate, i.e., a salt of lactic acid, which has to be recovered from the fermentation broth also containing proteins and lactic acid producing cells. In order to produce a more cost beneficial lactic acid and/or lactate, the fermentation has to utilise a micro-organism which produces a high concentration of lactate defined as overall concentration, but also defined as specific
25 productivity (Q_s), i.e., grams per gram of substrate, and volumetric productivity (Q_v), i.e., grams per litre of broth per hour.

Lactic acid is present in two enantiomeric forms, L-lactic acid, and D-lactic acid. As D-lactic acid is toxic to humans, and should not be present in food applications, there is a
30 demand for increased production of L-lactic acid. Lactic acid is produced during fermentation by means of an enzyme, lactate dehydrogenase, LDH. LDH is present in two forms, L-LDH for production of L-lactate, and D-LDH for the production of D-lactate.

The lactococci will thus also produce and activate a D-LDH, i.e., a lactate dehydrogenase producing the D-isomer.

US-A-4,885,247 relates to a method for recovery and purification of lactate salts from whole fermentation broth using electrodialysis, whereby the method is stated to efficiently recover the lactate as a concentrated liquid. The lactate recovered is then transformed into lactic acid.

Sjöberg, A., Persson, I., Quednau, M., and Hahn-Hägerdahl, B., in Applied Microbiology and Biotechnology 42(6):931-938, (1995) discuss the influence of limiting and non-limiting carbohydrate conditions of glucose and maltose metabolism in *Lactococcus lactis* spp. *lactis* strains. In this paper three different lactococci strains are shown with regard to their lactate production at growth under carbohydrate limiting and non-limiting conditions. The carbohydrates studied are maltose and glucose. One mutant, AS211 was formed from 19435 after a novobiocin-treatment. AS211 has a 20% higher production of lactate compared with the 19435 strain at continuous fermentation of glucose. AS211 was grown at $D = 0.6 \text{ h}^{-1}$.

STN database, CAPLUS, acc.no. 1992:549358, L-lactate production from xylose employing *Lactococcus lactis* IO-1, Biotechnology Letters, 14(7):599-604, (1992), and Dialog database, acc. no. 0137847, L-lactate production from xylose employing *Lactococcus lactis* IO-1-effect of inoculum C-source, xylose concentration, product inhibition and, mixed substrate on L-lactic acid production, Biotechnology Letters, 14(7):599-604, (1992) by Ishizaki, A. et al relates to work on *Lactococcus lactis* IO-1 with regard to different parameters which may influence the L-lactate production from a mixture of glucose and xylose. The basis for the study was to investigate whether L-lactate production may be optimised using a cheap raw material such as lignocellulose hydrolysate. Lignocellulose hydrolysate contains both xylose and glucose, which are used for the conversion into L-lactate. Using a mixture of glucose and xylose the production of L-lactate is 0.67 grams of lactate per gram of sugar. There is no indication in the paper of which subspecies (spp) *L. lactis* IO-1 belongs to.

Dialog database, acc. no. 07450183, General character and taxonomic studies of *Lactococcus lactis* IO-1 JCM 7638, Jour. of the faculty of agriculture, Kyushu

University, 35(1-2):1-8, (1990) by Ishazaki, A., et al, relates to the characterisation of the strain *L. lactis* IO-1, i.e., the strain discussed above. There is no indication in the paper of which subspecies (spp) *L. lactis* IO-1 belongs to.

5 Dialog database, acc. no. 00324956, Fermentative production of L-lactate from xylose, Conference paper, Developments in food engineering: Proceedings of the 6th International Congress on Engineered Food, Chiba, 2(4):552-554, (1993), by Ueda, T., et al, relates to *L. lactis* IO-1. Its ability of producing L-lactate from xylose is discussed.

10 STN database, CAPLUS, acc. no. 1994:571806, Cloning, sequencing, and comparison of three lactococcal L-lactate dehydrogenase genes, Microbiology 140(6):1301-1305, (1994), by Swindell, S. R., et al compares DNA sequences of genes coding for the enzyme L-lactate dehydrogenase (LDH) of different *Lactococcus* strains. It proves that the DNA sequences are strongly conserved between the different strains.

15

STN database, acc. no. 0167424, Stimulation of the rate of L-lactate fermentation using *Lactococcus lactis* IO-1 by periodic electrodialysis L-lactic acid and production, Jour. of Fermentation and Bioengineering, 77(5):508-512, (1994), by Vonktaveesuk, P., et al, discloses electrodialysis of the products obtained by fermentations with strain *L. lactis*

20 IO-1.

STN database, acc. no. 1991:523825, Differences between *Lactobacillus casei* spp *casei* 2206, and citrate-positive *Lactococcus lactis* spp. *lactis* 3022 in the characteristics of diacetyl production, Applied and Environmental Microbiology, 57(10):3040-3042,

25 (1991), by Kaneko, T., et al compares a lactococci, *Lactococcus lactis* spp. *lactis* and a lactobacilli, *Lactobacillus casei* spp. *casei* with regard to the production of diacetyl. In reality the LDH productions of the two strains are compared whereby the lactobacilli strain has 3 times higher production of LDH than the lactococci strain. The basic medium used is not disclosed.

30

Summary of the present invention

It has now been developed a new mutant of the wild type strain *L. lactis* spp. *lactis* 19435, which mutant strain, under given conditions produces high concentrations, expressed both

as Q_s and Q_v of L-lactate. Said mutant activates D-LDH, and L-LDH at different optima which provides for a differentiated production of D- and L-lactic acids.

The mutant can be used for high yield production of lactate to be used in food technology or as a commodity in chemical industry in general.

Detailed description of the invention

The present invention is based upon a mutant of *Lactococcus lactis* spp. *lactis* wild type strain 19435 (obtained from ATCC), which mutant has been deposited on the 4th of September, 2001 at Deutsche Sammlung von Mikroorganismen und Zellkulturen under deposition number DSM 14489 in accordance with the Budapest Treaty.

It is at present unknown where in the genetic code the mutation has occurred and to what extent the mutation has changed the genetic code.

At controlled, monitored fermentations (continuous fermentations) of the new mutant, herein called TMB5003, the lactate production has been analysed and calculations made show that TMB5003 has double the volumetric production compared to the wildtype strain 19435, and has a specific productivity which is 1.5 times that of the wildtype strain. The yield of lactate calculated as grams of lactate produced per gram of glucose added at the fermentation, was calculated to be the same for both strains at continuous fermentations.

The different tests made on the new mutant TMB5003 show, as will be evident from below that the new mutant has an unrestricted uptake of glucose which is transformed into lactate via a doubled metabolic capacity.

It has turned out that L-LDH, and D-LDH are produced or activated at different growth conditions, and thus the conditions can be chosen to produce optimal amounts of the lactate preferred, in this case the L-lactic acid or L-lactate.

Bacterial strains and cultivation conditions.

Escherichia coli DH5 α (Life Technologies Inc.) was grown in Luria-Bertani medium at 37 °C and erythromycin was added to a concentration of 250 $\mu\text{g ml}^{-1}$ when required. For standing batch cultures, *L. lactis* strains were cultivated at 30 °C in M17 medium (Oxoid) containing 10 g l $^{-1}$ sugar. For all lactococcal cultivations, sugars were autoclaved and
5 added separately to the cultures as well as erythromycin was added to a final concentration of 2 $\mu\text{g ml}^{-1}$ when required to select for TMB5003. In experiments using pH-controlled batch cultivations, the lactococcal strains were grown in a medium of the following composition (per litre): tryptone (Merck), 5 g; yeast extract (Merck), 5 g; casamino acids (Difco Laboratories), 1 g; K₂HPO₄, 2.5 g; KH₂PO₄, 2.5 g and MgSO₄·7
10 H₂O, 0.5 g, (pH 6.8). Carbohydrates were added to a final concentration of 10 g l $^{-1}$ respectively. The pH-controlled batch fermentations were performed at 30 °C in fermenters with a working volume of 800 ml. Stirring was set to 250 r.p.m. and the pH to 6.5, which was controlled by automatic base (3.0 M KOH) addition. The controlling device was a laboratory pH meter (Radiometer, Copenhagen, Denmark). The parent
15 cultures were grown overnight in the same media as the respective experimental cultures, in standing batch cultures at 30 °C. The inoculum 5% (vol/vol) was centrifuged, washed twice and resuspended in fresh culture medium without sugar, before being added to the experimental cultures.

20 For physiological characterisation and comparison of *L. lactis* spp. *lactis* ATCC19435 and *L. lactis* spp. *lactis* TMB5003, these strains were cultivated in semi-defined medium (SD3) according to van Niel and Hahn-Hägerdahl, (1999). All components except for potassium phosphates and water, were sterile filtered. Glucose was autoclaved and added separately to the medium at a final concentration of 5 g l $^{-1}$. Continuous cultivations were
25 performed using chemostat conditions in Biostat[®] A fermenters (B. Braun Biotech International, Germany). The volume in the fermenters was kept at 1 l. The temperature was set at 30 °C and the pH was maintained at 6.5 using automatic base (10 M KOH) addition by the use of an automatic controlling device, micro-DCU system (B. Braun Biotech International, Germany). The stirring was kept at 150 r.p.m. using a MCU-200
30 system (B. Braun Biotech International, Germany). Anaerobic conditions were withheld by continuous nitrogen flushing through the medium at 0.2 ml min $^{-1}$. Precultures for fermentations were grown overnight in M17 medium containing 10 g l $^{-1}$ glucose. Precultures were harvested by centrifugation at 5 000×g, 2 °C for 10 min, washed twice

and resuspended in sterile double distilled water and finally inoculated at 2.5 % (vol/vol) into SD3 medium containing 5 g l⁻¹ glucose. Continuous cultivations were started when the glucose was depleted from the batch cultures. Continuous fermentations were run in duplicates at each dilution rate for both lactococcal strains.

5

Measurement of growth, substrate consumption and product formation.

Measuring the optical density (OD) at 620 nm, using appropriate dilutions, on a Hitachi U-2000 spectrophotometer (Hitachi Ltd., Tokyo, Japan), monitored cell growth. Dry weight was measured for all cultures. Samples for substrate and product determination were filtered immediately through 0.2 µm filters after sampling and kept at -20°C until analysis. Glucose, lactate, formate, acetate and ethanol were separated at 45 °C on an ionexchange column (Aminex HPX-87H, BioRad) and quantified using a refractive index detector (Shimadzu, Japan). The mobile phase was 5 mM H₂SO₄ at a flow rate of 0.6 ml min⁻¹.

15

Cell extract preparation, protein determination and enzyme assays.

Lactococcal cells were withdrawn from the cultivations at appropriate times and harvested by centrifugation at 5 000×g, 2 °C for 10 min. The cells were washed twice and resuspended in 20 mM triethanolamine buffer, pH 7.2, containing 0.5 mM EDTA and 0.5 mM dithiotreitol. Disintegration of cells was performed by vortexing (3×5 min) at 8 °C by the use of glass beads (0.5 mm, KEBO). Cell debris was removed by centrifugation at 19,500×g, 2 °C, for 15 min. Cell extracts were kept at -80 °C until used. The protein concentration was determined according to the method of Bradford [Bradford, (1976)]. Bovine serum albumin was used as a standard.

25

All chemicals used in the enzyme assays were obtained from Sigma-Aldrich. LDH and PK activity was measured in the direction of NADH oxidation at 340 nm according to Hillier and Jaga, (1982), and Crow and Pritchard, (1982), respectively. The activity of PFK and GAPDH was determined according to the methods of Plaxton and Storey, (1986) and Brooks and Storey, (1988), respectively.

30

Glucose transport measurement.

The uptake of glucose by lactococcal cells was measured according to the zero-*trans*-influx assay adapted for bacterial cells. Cells were collected at appropriate times and harvested by centrifugation at 5 000×g, 2 °C, for 10 min. Cell pellets were washed twice in ice-cold 0.1 M potassium phosphate buffer, pH 6.5, and finally resuspended to a dry weight of 25-30 mg ml⁻¹ using the same buffer. Cells were kept on ice until used. In the assay 20 µl of 0.1 M potassium phosphate buffer, pH 6.5, and 20 µl of cell resuspension were added to a plastic 5 ml reaction vial (Sarstedt) and incubated at 30 °C for 5 min. Ten microlitres of ¹⁴C-labelled glucose (Amersham Life Science) of concentrations between 0.3125 mM to 200 mM was added to the mixture to a final specific activity of 200 counts per minute (cpm) per nmol, and the assay was started by vortexing the vial shortly. The assay was allowed to proceed for 10 seconds, measured by the use of a metronome, and stopped by adding 3 ml of ice cold 0.5 M glucose from a dispenser. The reaction mixture was rapidly filtered through a glass micro-fibre filter (GF/F, 25 mm, Merck Eurolab). The reaction vial was washed twice with 3 ml ice-cold 0.5 M glucose solution and finally the filter equipment was also washed twice with ice-cold 0.5 M glucose solution. The filter was placed in a scintillation vial containing 5 ml scintillation solution (Ecoscint™ A, Hinzte AB, Sweden). Assays were run in duplicates and for every cell resuspension and for every glucose concentration a background sample was prepared. These samples were prepared and handled as the other tests but the assay was not started by vortexing, instead 3 ml ice-cold 0.5 M glucose solution was dispensed into the vial and the reaction mixture was filtered immediately.

Genetic techniques and development of *L. lactis* TMB5003.

All DNA-modifying enzymes were obtained from Roche Diagnostics Scandinavia AB, Sweden. Plasmid preparations were performed using a Bio-Rad Quantum prep kit (Bio-Rad) and chromosomal DNA was prepared using an Easy-DNA™ kit (Invitrogen). Polymerase chain reaction (PCR) was performed using *Pwo*II polymerase according to manufacturer's description. DNA fragments were purified from agarose gel using a Qiaquick kit (Qiagen). Restriction enzyme digestions and ligations were performed according to standard procedures [Sambrook, (1989)]. Ultra-competent *E. coli* cells were prepared and transformed as previously described [Inoue, (1990)]. Preparation and transformation of lactococcal cells were performed according to a protocol by Holo and Nes, (1989).

A 850 bp internal fragment of the *L. lactis* maltose phosphorylase encoding gene, *malP*, [Nilsson, Microbiology, 147:1565-1573, (2001)] was amplified by PCR using primers 5'-ggcggatcctaaaggatttactgg-3' (forward), containing a *Bam*HI restriction enzyme recognition site and 5'-ggcctgcagcaacttctcgcttg-3' (reverse) containing a *Pst*I restriction recognition site and *L. lactis* ssp. *lactis* 19435 chromosomal DNA as template. The PCR product was cleaved with restriction enzymes *Bam*HI and *Rsa*I, resulting in a 450 bp product. A minimal integration vector, pFL20, not able to replicate in lactococci, developed by Levander et al., (2001), was digested with suitable restriction enzymes and ligated with the 450 bp *malP* internal fragment. The resulting construct, denoted pTMB5003, was propagated in *E. coli* and further transformed into *L. lactis* ssp. *lactis* 19435. Four transformants were obtained on erythromycin selective plates due to a single cross-over event in the *malP* of *L. lactis*. All transformants appeared to have the same growth behaviour in glucose and maltose cultivations, respectively. One transformant, called *L. lactis* ssp. *lactis* TMB5003, was chosen for further investigations.

The new mutant has been tested during glucose fermentation runs, and thereby been compared with the type strain.

Growth and product formation.

L. lactis ssp. *lactis* 19435 and *L. lactis* ssp. *lactis* TMB5003 were grown in pH-controlled batch cultures to assess and compare their growth behaviour on glucose (Fig. 2). The maximum specific growth rate of TMB5003 was twice the one that was determined for wild-type lactococci under the same growth conditions. When measuring the consumption of glucose during the cultivations it was obvious that TMB5003 consumed glucose approximately twice as fast as the wild type. Batch cultivations using lactose or maltose as sole carbon source resulted in no growth for TMB5003 in any of the cultivations while 19435 grew with the same specific growth rate on lactose as in glucose cultivations and with a slightly lower rate on maltose (data not shown). The inability of TMB5003 to ferment lactose was confirmed by investigating the plasmid content of these cells. TMB5003 had lost one plasmid, compared to wild-type lactococci, most likely to be the one harbouring the *lac*-operon (data not shown) [de Vos, (1990); de Vos, (1989); Maeda, (1986)]. The inability of the mutant strain to ferment maltose was due to the fact that the maltose operon was disrupted by insertion of pTMB5003 in the maltose

phosphorylase encoding gene, *malP* (Nilsson and Rådström, (2001)). It is believed that the alteration in the maltose operon does not promote the effect on glucose metabolism in TMB5003 and therefore this is not further discussed in the current presentation.

- 5 Product formation was investigated when lactococcal cells were cultivated continuously using glucose as sole carbon source under limiting and non-limiting conditions. TMB5003 was shown to produce lactate by a volumetric productivity twice as high as that for 19435 under glucose non-limiting conditions (Table 1). However, when glucose was limiting, the lactate productivity decreased markedly for TMB5003, beyond the
10 values of wild-type lactococci. These results suggest that the mutant strain has a stronger demand for excess glucose, in order to produce lactate as main fermentation end product, than the wild type strain. It is known from earlier studies that a shift towards mixed acid product formation occurs when glucose becomes limited in a lactococcal cultivation or when certain other sugars, such as maltose, are used as sole carbon sources [Lohmeier-
15 Vogel, (1986); Coccagn-Bousquet, (1996); Sjöberg, (1995)]. This is due to a diminished flux through glycolysis and a lower NADH/NAD⁺ ratio. It has further been explained that the bottleneck under such conditions most probably is at the level of sugar transport.

Glucose uptake and enzyme activities.

- 20 Measurements of glucose uptake in lactococcal cells cultivated under non-limiting glucose conditions were enabled using ¹⁴C-labelled glucose. The trend in glucose transport, using a range of glucose concentrations of 0.3125 mM to 200 mM, was shown to be saturating when higher glucose concentrations than approximately 20 mM was used in the assay for 19435 (Fig. 1). On the other hand, when TMB5003 was applied in the
25 same assay the trend of glucose uptake did not show a saturating curve, although, for which reason is to us unknown, lower values of specific uptake rates were obtained. These results indicate that a regulatory function aimed at controlling the influx of sugar into lactococci is affected in the mutant cells. This may also be reflected upon when considering the more pronounced effect of shift towards mixed acid product formation for
30 the mutant lactococci strain at glucose levels not considered to be limiting for the wild type lactococci (Table 1). There are evidences of two glucose transport systems in *L. lactis*, PTS and permease mediated, respectively [Thompson, (1985); Thompson, (1983)]. Due to the fact that glucose is transported by the mannose specific PTS in *L. lactis* it was tempting to check the growth rate of the lactococcal strains on this carbohydrate. Hence,

both strains were shown to grow at the same specific growth rate as in their glucose cultures, respectively (data not shown). Despite the fact that many reports describe the regulatory role of PTS on the uptake of PTS- and non-PTS-sugars, via the glucose effect, there are not much information concerning regulation of uptake of glucose itself. The heat-stable protein (HPr) of the PTS and the HPr kinase are known to take part in the inducer expulsion and exclusion phenomena, and there are also suggestions of these to play a role in control of glucose uptake [Cocaign-Bousquet, (1996); Saier Jr., (1996)]. It is tempting to speculate that these proteins might be affected in TMB5003 and thus mediate the altered mode of glucose uptake in this strain. However, due to the fact that TMB5003 did also have an improved specific growth rate when cultivated on mannose, but not on trehalose, also suggested to be transported by PTS (Nilsson and Rådström, (2001)), further speculations occur concerning the mannose specific components of the mannose/glucose uptake system to be affected in the mutant strain.

In *L. lactis* phosphofructokinase (PFK), pyruvate kinase (PK) and lactate dehydrogenase (LDH) are the key enzymes in the central pathway of energy production, the conversion of carbohydrates into lactic acid [Llanos, (1993)]. The genes encoding these enzymes are located together in an operon, denoted as the *las*-operon, on the chromosome. Recent results have been obtained concerning the role of PFK on glycolytic flux in *L. lactis* [Andersen, (2001)]. The conclusions were that glycolytic and lactate fluxes were decreased proportionally by a twofold reduction of PFK activity. Activities of the enzymes encoded by the *las*-operon were investigated in 19435 and TMB5003 cultivated at different dilution rates (Table 2). For mutant lactococci no difference in activities of PK and PFK could be detected in cell extracts from cultivations at the three different dilution rates. However, activities of LDH detected in TMB5003 cells collected at different growth rates differed markedly. LDH activity was higher in fast growing cells and it was showed to be twenty times higher in TMB5003 than in 19435 compared at the highest dilution rate for each strain respectively. These results are in consistence with recent findings of high LDH activities at high glycolytic rates for lactococci [Even, (2001)]. The markedly higher LDH activity in TMB5003 is most probably a response to the altered transport of glucose into the cells. Since the lactococci are provided with excess glucose and glycolytic flux is high, the mode of pyruvate conversion has to proceed accordingly due to the requirement of regeneration of NAD^+ . This is performed by the reduction of pyruvate to lactate catalysed by LDH, which activity is therefore

required to be kept high. In *L. lactis* it has been shown that *in vivo* glyceraldehyde phosphate dehydrogenase (GAPDH) activity limits the glycolytic flux on rapidly metabolizable sugars, due to the inhibition of a high NADH/NAD⁺ ratio [Even, (1999)]. Therefore, we wanted to investigate if TMB5003 possessed an altered activity of GAPDH compared to wildtype lactococci. However, no difference in activities could be detected in the strains (Table 2).

The investigations of glucose uptake of the respective strain at controlled fermentation runs, at highest possible dilution rate with regard to each individual strains, were carried out. The results show that the new mutant TMB5003 has an unlimited uptake of glucose while a certain saturation of the uptake of glucose by the wildtype strain 19435 is noted. This is evident from FIG. 1, which shows specific uptake rates given as nmol/min/mg of cells added to the assay, versus the concentration of glucose given as mM added to the assay.

The concentrations of glucose were varied between 0 to 80 mM, and the specific uptake rate varied between 10 to 150 nmol/min/mg cells added, with regard to the wildtype strain, and between 5 to 55 nmol/min/mg cells added with regard to the new mutant.

The fermentation rates when grown on a glucose medium were compared as well, as evident from FIG. 2. The new mutant TMB5003 ferments glucose at the double rate compared to the wildtype strain. FIG. 2 shows both growth on glucose (optical density graph, OD-curve), and glucose consumption of the wildtype strain 19435, and TMB5003, respectively.

In Table 1 below the results of studied lactate formation is given.

TABLE 1

	<i>L. lactis</i> ATCC 19435			<i>L. lactis</i> TMB5003		
Dilution rate (D)	0.1	0.2	0.4	0.1	0.4	0.8
Y _{LA/tot}	0.7	0.6	0.8	0.1	0.3	0.8
Q _s (g/g/h)	0.4	0.6	2.0	0.03	0.5	3.1
Q _v (g/l/h)	0.3	0.6	1.6	0.04	0.6	3.4

10 $Y_{LA/tot}$ stands for yield of lactic acid in relation to total production of fermentation products.

Q_s is specific productivity, i.e., grams per gram of substrate.

Q_v is volumetric productivity, i.e., grams per litre of broth per hour.

15

Furthermore the production of different enzymes present was analysed. The enzymes of interest are hereby the LDH, phosphofructokinase (PFK), the pyruvate kinase (PK) and the glyceraldehyde phosphate dehydrogenase (GAPDH). Comparisons are made at the highest D of the respective strain. The results are given in Table 2 below.

20

TABLE 2

Enzyme	<i>L. lactis</i> 19435	<i>L. lactis</i> TMB5003
	D = 0.4	D = 0.8
	U/mg	U/mg
25 Lactate dehydrogenase (LDH)	0.35	7.5
Phosphofructokinase (PFK)	0.5	0.5
Pyruvate kinase (PK)	0.15	0.1
30 Glyceraldehydephosphate dehydrogenase (GAPDH)	0.03	0.02

As evident from Table 2 above the production and activation of LDH is outstanding as compared to the wildtype, as it is at least 20 times that of the wildtype, which means that the present new strain has a higher capacity of producing lactate into the growth medium. With regard to the other enzymes analysed there are no differences to be seen between the two strains.

The L-lactate produced and recovered can be used to adjust pH in food, as a taste enhancer in food, as well as a preservative of food, whereby the preservative effect is due to a lowering of pH, as well as due to the weak acid itself, which prevents growth of a number of micro-organisms in food and feedstuff. Accumulation of anions intracellularly or uncoupling of ATP-synthase is thereby the most probable mechanisms of growth inhibition.

Further, the lactate can be used for the treatment of paper and metallic surfaces.

The lactate can be used for a polymerisation into poly-lactic acid, PLA, which is a biodegradable polymer. Further, the lactate can be used in the production of other compounds, such as propylene glycol, propylene oxide, acetaldehyde, ethanol, acrylates, and acrylic esters.

PLA as such can be applied in medical applications in the form of implants and sutures, production of items used for controlled release of drugs, and pesticides. The polylactate can be used in the manufacture of package materials, as well as biodegradable disposable items.

The present strain can be grown in different media, such as complex media based on tryptone, yeast extracts, and casamino acids. Glucose can be added as an external carbon source. However, complex media are not preferred, as the amount of sugar (glucose) which results in lactate formation shall be controlled. In tryptone and yeast extracts there will be unknown components present which may make this difficult. A semi-defined media, such as SD3 and an addition of glucose up to 5 g/l can be used.

The semi-defined medium (E.W.J van Niel and B. Hahn-Hägerdal (1999), "Nutritient requirements of lactococci in defined growth media", Applied Microbial Biotechnology 52:617-627) used for cultivating the present strain TMB5003 in the investigations made is composed of

5 SD3 medium (per litre)

	Casamino acids	10 g
	K ₂ HPO ₄	2,5 g
	KH ₂ PO ₄	2,5 g
	MgSO ₄ · 7H ₂ O	0.5 g
10	Yeast nitrogen base	5 g
	(except casamino acids (Difco))	
	Asparagine	0.4 g
	Reduced glutathione	10 mg
	Uracil	60 mg
15	Adenine	30 mg
	Guanine	30 mg
	Vitamin solution	10 ml
	Trace element solution	1 ml
	Glucose	5 g

20

Vitamin solution (per litre)

	D-biotin	10 mg
	Pyridoxal-HCl	206 mg
	Folic acid	100 mg
25	Riboflavin	100 mg
	Niacinamide	100 mg
	Thiamine-HCl	100 mg
	Ca-D-panthotenate	95 mg
	p-Aminobenzoic acid	10 mg

30

Trace element solution (per litre)

	Ca ₂ -EDTA	15 g
	ZnSO ₄ · 7H ₂ O	4.5 g
	MnCl ₂ · 2H ₂ O	1 g
5	CoCl ₂ · 6H ₂ O	0.3 g
	CuSO ₄ · 5H ₂ O	0.3 g
	Na ₂ MoO ₄ · H ₂ O	0.4 g
	CaCl ₂ · 2H ₂ O	4.5 g
	FeSO ₄ · 7H ₂ O	3 g
10	H ₃ BO ₃	1 g
	KI	0.1 g

The solutions of vitamins, trace elements, nucleic acid bases, yeast nitrogen base, asparagine and reduced glutathione were filter sterilised and added to the medium aseptically. The other components
15 were autoclaved separately. At selection of TMB5003 2µg of erythromycin /ml medium is added.

It is preferred that the present mutant TMB5003 is grown at high dilution rate, i.e., with a complete addition of glucose, and absolutely no restricted addition thereof to obtain and maintain a maximal lactate production. Thus a dilution rate of at least 0.5h⁻¹, preferably at
20 least 0.7h⁻¹, and most preferably at least 0.8h⁻¹ is used. At a restricted addition of glucose during growth the lactate production tends to drop to the benefit of formation of by-products such as other acids such as acetate (acetic acid) and formate (formic acid).

It is, so far, very probable that the limiting factor of the wildtype strain to produce lactate
25 is the ability of the strain to transport glucose. The flux by means of the glucolysis is probably not limiting. The final pyruvate metabolism may be limiting. LDH is the enzyme which transforms pyruvate into lactate and this enzyme has turned out to have different temperature optima depending on which strain of lactococci that is used at the fermentation of glucose. pH, as well, may have an effect upon the activity of LDH.
30 Furthermore, the redox balance influences the efficiency of the LDH. The presence of the co-factors NAD⁺/NADH is the controlling factor.

The conditions for enhancing L-LDH activity are fermentation in a medium having a pH above 6, and a temperature of up to 30°C, at which conditions L-lactate is substantially

the only isomer produced, while D-LDH is activated at pH 4-5, and 33.5 to 40°C. Thus the fermentation of the present mutant should be carried out at such conditions that enhance activity of L-LDH to enhance formation of L-lactate. Thus, the pH and temperature conditions for production of L-lactate are pH 6 to 7, preferably 6.0 to 6.5, and using a temperature of 25 to 30°C, preferably 27.5 to 30°C.

FIGURE LEGENDS

FIG. 1A shows the specific glucose uptake rate *Lactococcus lactis* spp. *lactis* 19435 cultivated at a dilution rate of 0.4 h^{-1} .

FIG. 1B shows the specific glucose uptake rate by *Lactococcus lactis* spp. *lactis*

5 TMB5003 cultivated at a dilution rate of 0.8 h^{-1} .

FIG. 2. shows the glucose consumption versus time, and the change of optical density (OD) versus time.

CLAIMS

1. A mutant of *Lactococcus lactis* spp. *lactis*, which produces lactate at high volumetric and specific productivity as well as high amounts of lactate dehydrogenase.
- 5 2. A mutant according to claim 1, which has a volumetric productivity of at least twice that of the *Lactococcus lactis* spp. *lactis* 19435.
3. A mutant according to claim 1, which has a specific productivity of at least 1.5 times that of the *Lactococcus lactis* spp. *lactis* 19435.
- 10 4. A mutant according to claim 1, which has a production of lactate dehydrogenase of at least twenty times that of the *Lactococcus lactis* spp. *lactis* 19435.
5. A mutant according to claims 1-4, denoted TMB5003, deposited at Deutsche
15 Sammlung von Microorganismen und Zellkulturen under deposition number DSM 14489.
6. The use of *Lactococcus lactis* spp. *lactis* TMB5003 in the production of lactate.
7. The use of *Lactococcus lactis* spp. *lactis* TMB5003 for the production of the enzyme
20 lactate dehydrogenase, LDH, in particular L-lactate dehydrogenase, L-LDH.
8. Method for the production of lactate on a glucose containing medium, wherein an inoculum of *Lactococcus lactis* spp. *lactis* TMB5003 is grown on a medium comprising glucose as carbon source.
- 25 9. Method according to claim 8, wherein the growth is carried at conditions optimised for production of L-lactate.
10. Method according to claims 8-9, wherein the growth is continuous at a dilution rate of
30 at least 0.5 h^{-1} .
11. Method according to claim 10, wherein the growth is continuous at a dilution rate of at least 0.7 h^{-1} .

12. Method according to claim 11, wherein the growth is continuous at a dilution rate of at least 0.8 h^{-1} .

5 13. Method according to claims 8-12, wherein the growth is carried out at unrestricted feed of glucose.

14. Method according to claims 8-13, wherein the growth is carried out at a pH of above 6, preferably pH 6-7.

10 15. Method according to claims 8-14, wherein the growth is carried out at a temperature of between 25 and 30 °C, preferably 27.5 to 30°C.

16. The use of lactate produced by fermenting the mutant *Lactococcus lactis* spp. *lactis* TMB5003 in food and as a chemical commodity in general.

1/2




FIG. 1A




FIG. 1B

2/2




FIG.2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 02/01805

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C12N 1/20, C12P 7/56 // (C12N 1/20, C12R 1:225)
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C12N, C12R

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, BIOSIS, MEDLINE, CHEM.ABS.DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Appl. Microbiol Biotechnol, Volume 42, 1995, A. Sjöberg et al: "The influence of limiting and non-limiting growth conditions on glucose and maltose metabolism in Lactococcus lactis ssp. lactis strains", pages 931-938, page 932, column 1, figures 285 --	1-6,8-15
X	FEMS Microbiology Letters, Volume 107, 1993, Polly K. Dinsmore et al: "Insertional mutagenesis in Lactococcus lactis subsp. lactis mediated by IS946", pages 43-48, abstract, page 46, column 1 - page 47, column 1 --	1-5

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 January 2003

Date of mailing of the international search report

13-01-2003

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

TERESE PERSSON/BS

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 02/01805

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FEMS Microbiology Letters, Volume 81, 1991, Colin Hill et al: "Isolation of chromosomal mutations of Lactococcus lactis subsp. lactis biovar. diacetylactis 18-16 after introduction of Tn919", pages 135-140, summary; page 138 --	1-5
X	Journal of Bacteriology, Volume 177, No. 24, 1995, Jean Law et al: "A System To Generate Chromosomal Mutations in Lactococcus lactis Which Allows Fast Analysis of Targeted Genes", pages 7011-7018, abstract --	1-5
X	WO 9713842 A1 (MIDWEST RESEARCH INSTITUTE), 17 April 1997 (17.04.97), page 1, line 16 - line 17 --	16
A	Applied and Environmental Microbiology, Volume 67, No. 10, 2001, Fredrik Levander et al: "Physiological Role of Beta-Phosphoglucomutase in Lactococcus lactis", pages 4546-4553 --	1-16
A	Current Opinion in Biotechnology, Volume 10, 1999, Jeroen Hugenholtz et al: "Metabolic engineering of lactic acid bacteria: overview of the approaches and results of pathway rerouting involved in food fermentations", pages 492-497 --	1-16
A	Microbiology, Volume 147, 2001, Ulrika Nilsson et al: "Genetic localization and regulation of the maltose phosphorylase gene, malP, in Lactococcus lactis", pages 1565-1573 -- -----	1-16

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE02/01805

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-16 (partially)
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

see next sheet

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE02/01804

Present claims 1-5 relate to a mutant of *Lactococcus lactis* spp. *lactis* defined by reference to a number of desirable characteristics and properties, namely a high volumetric and specific productivity of lactate production as well as a high production of lactate dehydrogenase. The claims cover all mutants of *Lactococcus lactis* spp. *lactis* having these characteristics and properties, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for only a very limited number of such mutated *Lactococcus lactis* spp. *lactis*. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Independent of the above reasoning, the claims also lacks clarity (Article 6 PCT). An attempt is made to define the mutated *Lactococcus lactis* spp. *lactis* by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

The present description does not disclose where in the genetic code the mutation, which is responsible for the features of the mutant, has occurred or how the mutation has changed the genetic code (see page 5, line 13-14). As a result, the search has mainly been focused on mutated *Lactococcus lactis* spp. *lactis*, in which the gene that encodes the maltose phosphorylase has been mutated. This is the only genetic feature that the mutant in the description has been characterised to have. A search has also been carried out for mutated *Lactococcus lactis* spp. *lactis* which have been characterised to have the features described above (i.e. high lactate and lactate dehydrogenase production).

As a consequence, additional claims relating to this mutant *Lactococcus lactis* spp. *lactis* have been searched with the same limitations as described above.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established will not be the subject of an international preliminary examination (Rule 66.1(e) PCT). This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 02/01805

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9713842	A1	17/04/97	AU	697432 B	08/10/98
				AU	7373896 A	30/04/97
				CA	2234170 A	17/04/97
				EP	0859829 A	26/08/98
				NO	981580 A	09/06/98
				US	5798237 A	25/08/98
